

Clinical Policy: Icosapent Ethyl (Vascepa)

Reference Number: ERX.NPA.140

Effective Date: 03.01.20

Last Review Date: 02.21

Line of Business: Commercial, Medicaid

[Revision Log](#)

See **Important Reminder** at the end of this policy for important regulatory and legal information.

Description

Icosapent ethyl (Vascepa[®]) is an ethyl ester of eicosapentaenoic acid (EPA).

FDA Approved Indication(s)

Vascepa is indicated:

- As adjunct to maximally tolerated statin therapy to reduce the risk of myocardial infarction, stroke, coronary revascularization, and unstable angina requiring hospitalization in adult patients with elevated triglyceride (TG) levels (≥ 150 mg/dL) and:
 - Established cardiovascular disease (CVD) or
 - Diabetes mellitus and 2 or more additional risk factors for cardiovascular disease
- As an adjunct to diet to reduce TG levels in adult patients with severe (≥ 500 mg/dL) hypertriglyceridemia

Limitation(s) of use: The effect of Vascepa on the risk for pancreatitis in patients with severe hypertriglyceridemia has not been determined.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

Health plan approved formularies should be reviewed for all coverage determinations. Requirements to use preferred alternative agents apply only when such requirements align with the health plan approved formulary.

It is the policy of health plans affiliated with Envolve Pharmacy Solutions[™] that Vascepa is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Hypertriglyceridemia without ASCVD (must meet all):

1. Diagnosis of hypertriglyceridemia;
2. Age ≥ 18 years;
3. Fasting TG ≥ 500 mg/dL (lab must be dated within 90 days);
4. Failure of a ≥ 3 consecutive month trial of fibrates therapy in the last 6 months at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
5. Failure of omega-3-acid ethyl esters (generic Lovaza[®]) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
6. Dose does not exceed 4 g (4 capsules) per day.

Approval duration: 6 months

B. Reduction of Cardiovascular Disease Risk (must meet all):

1. Diagnosis of one of the following (a or b):
 - a. Atherosclerotic cardiovascular disease (ASCVD) as evidenced by a history of any one of the following conditions (i-vii);
 - i. Acute coronary syndromes;

- ii. Clinically significant coronary heart disease (CHD) diagnosed by invasive or noninvasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography, or nuclear imaging);
 - iii. Coronary or other arterial revascularization;
 - iv. Myocardial infarction;
 - v. Peripheral arterial disease presumed to be of atherosclerotic origin;
 - vi. Stable or unstable angina;
 - vii. Stroke or transient ischemic attack (TIA);
 - b. Diabetes with ≥ 2 CVD risk factors (*see Appendix G*);
2. Age ≥ 45 years;
3. Documentation (labs must be dated within 90 days) of fasting TG between 150-499 mg/dL;
4. Documentation of low-density lipoprotein cholesterol (LDL-C) (labs must be dated within 90 days) between 41-100 mg/dL;
5. For members on statin therapy, both of the following (a and b):
 - a. Vascepa is prescribed in conjunction with a statin at the maximally tolerated dose;
 - b. Member has been adherent for at least the last 4 months to maximally tolerated doses of one of the following statin regimens (i, ii, or iii):
 - i. A high intensity statin (*see Appendix D*);
 - ii. A moderate intensity statin (*see Appendix D*), and member has one of the following (a or b):
 - a) Intolerance to two high intensity statins;
 - b) A statin risk factor (*see Appendix E*);
 - iii. A low intensity statin, and member has one of the following (a or b):
 - a) Intolerance to one high and one moderate intensity statins;
 - b) A statin risk factor (*see Appendix E*) and history of intolerance to two moderate intensity statins;
6. For members not on statin therapy, member meets one of the following (a or b):
 - a. Statin therapy is contraindicated per Appendix F;
 - b. For members who are statin intolerant, member has tried at least two statins, one of which must be hydrophilic (pravastatin, fluvastatin, or rosuvastatin), and member meets one of the following (i or ii):
 - i. Member has documented statin risk factors (*see Appendix E*);
 - ii. Member is statin intolerant due to statin-associated muscle symptoms (SAMS) and meets both of the following (a and b):
 - a) Documentation of intolerable SAMS persisting at least two weeks, which disappeared with discontinuing the statin therapy and recurred with a statin re-challenge;
 - b) Documentation of re-challenge with titration from lowest possible dose and/or intermittent dosing frequency (e.g., 1 to 3 times weekly);
7. Dose does not exceed 4 g (4 capsules) per day.

Approval duration: 6 months

C. Other diagnoses/indications

1. Refer to ERX.PA.01 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

II. Continued Therapy

A. Hypertriglyceridemia without ASCVD (must meet all):

1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by one of the following (a or b):
 - a. Initial re-authorization: 20% reduction in TG levels from baseline;
 - b. Subsequent re-authorizations: continued reduction or maintenance in reduction of TG levels from baseline;
3. If request is for a dose increase, new dose does not exceed 4 g (4 capsules) per day.

Approval duration: 12 months

B. Reduction of Cardiovascular Disease Risk (must meet all):

1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by no increase in TG and LDL-C levels from baseline;
3. If statin tolerant, documentation of adherence to a statin at the maximally tolerated dose;
4. If request is for a dose increase, new dose does not exceed 4 g (4 capsules) per day.

Approval duration: 12 months

C. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions and documentation supports positive response to therapy.

Approval duration: Duration of request or 12 months (whichever is less); or

2. Refer to ERX.PA.01 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – ERX.PA.01 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ASCVD: atherosclerotic cardiovascular disease

CVD: cardiovascular disease

EPA: eicosapentaenoic acid

FDA: Food and Drug Administration

LDL-C: low-density lipoprotein cholesterol

SAMS: statin-associated muscle symptoms

TG: triglyceride

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
fenofibrate (TriCor®)	HyperTG without ASCVD: 48-145 mg PO QD	145 mg/day
gemfibrozil (Lopid®)	HyperTG without ASCVD: 600 mg PO BID	1,200 mg/day
omega-3-acid ethyl esters (Lovaza®)	HyperTG without ASCVD: 4 g PO QD or 2 g PO BID	4 g/day

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): known hypersensitivity (e.g., anaphylactic reaction) to Vascepa or any of its components
- Boxed warning(s): none reported

Appendix D: High and Moderate Intensity Daily Statin Therapy for Adults

High Intensity Statin Therapy <i>Daily dose shown to lower LDL-C, on average, by approximately ≥ 50%</i>
<ul style="list-style-type: none"> • Atorvastatin 40-80 mg • Rosuvastatin 20-40 mg
Moderate Intensity Statin Therapy <i>Daily dose shown to lower LDL-C, on average, by approximately 30% to 50%</i>
<ul style="list-style-type: none"> • Atorvastatin 10-20 mg • Fluvastatin XL 80 mg

<p>Moderate Intensity Statin Therapy <i>Daily dose shown to lower LDL-C, on average, by approximately 30% to 50%</i></p> <ul style="list-style-type: none"> • Fluvastatin 40 mg BID • Lovastatin 40 mg • Pitavastatin 1-4 mg • Pravastatin 40-80 mg • Rosuvastatin 5-10 mg • Simvastatin 20-40 mg
<p>Low Intensity Statin Therapy <i>Daily dose shown to lower LDL-C, on average, by <30%</i></p> <ul style="list-style-type: none"> • Simvastatin 10 mg • Pravastatin 10-20 mg • Lovastatin 20 mg • Fluvastatin 20-40 mg

Appendix E: Statin Risk Factors

<p>Statin Risk Factors</p> <ul style="list-style-type: none"> • Multiple or serious comorbidities, including impaired renal or hepatic function • Unexplained alanine transaminase (ALT) elevations > 3 times upper limit of normal, or active liver disease • Concomitant use of drugs adversely affecting statin metabolism • Age > 75 years, or history of hemorrhagic stroke • Asian ancestry

Appendix F: Statin Contraindications

<p>Statins</p> <ul style="list-style-type: none"> • Decompensated liver disease (development of jaundice, ascites, variceal bleeding, encephalopathy) • Laboratory-confirmed acute liver injury or rhabdomyolysis resulting from statin treatment • Pregnancy, actively trying to become pregnant, or nursingImmune-mediated hypersensitivity to the HMG-CoA reductase inhibitor drug class (statins) as evidenced by an allergic reaction occurring with at least TWO different statins
--

Appendix G: Risk Factors for CVD (2020 ADA and 2013 AHA Guidelines)

- Men and women ≥ 50 years of age
- Cigarette smoker or stopped smoking within 3 months
- Hypertension (blood pressure ≥ 140 mmHg systolic OR ≥ 90 mmHg diastolic) or on antihypertensive medication
- HDL-C ≤ 40 mg/dL for men or ≤ 50 mg/dL for women
- Renal dysfunction: creatinine clearance (CrCL) > 30 and < 60 mL/min or chronic kidney disease
- Micro- or macroalbuminuria
 - Microalbuminuria: either a positive micral or other strip test (may be obtained from medical records), an albumin/creatinine ratio ≥ 2.5 mg/mmol, or an albumin excretion rate on timed collection ≥ 20 mg/min all on at least two successive occasions
 - Macroalbuminuria: Albustix or other dipstick evidence of gross proteinuria, an albumin/creatinine ratio ≥ 25 mg/mmol, or an albumin excretion rate on timed collection ≥ 200 mg/min all on at least two successive occasions
- Obesity/overweight
- Dyslipidemia
- Family history of premature coronary disease

Appendix H: General Information

- The diagnosis of SAMS is often made based on clinical criteria. Typical SAMS include muscle pain and aching (myalgia), cramps, and weakness. Symptoms are usually bilateral and involve

large muscle groups, including the thigh, buttock, back, and shoulder girdle musculature. In contrast, cramping is usually unilateral and may involve small muscles of the hands and feet. Symptoms may be more frequent in physically active patients. Symptoms often appear early after starting statin therapy or after an increase in dose and usually resolve or start to dissipate within weeks after cessation of therapy, although it may take several months for symptoms to resolve completely. Persistence of symptoms for more than 2 months after drug cessation should prompt a search for other causes or for underlying muscle disease possibly provoked by statin therapy. The reappearance of symptoms with statin rechallenge and their disappearance with drug cessation offers the best evidence that the symptoms are truly SAMS.

- Pravastatin, fluvastatin, and rosuvastatin are hydrophilic statins which have been reported to confer fewer adverse drug reactions than lipophilic statins.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Hypertriglyceridemia without ASCVD	2 g PO BID	4 g/day
Reduction of CVD risk		

VI. Product Availability

Capsules: 0.5 g, 1 g

VII. References

1. Vascepa Prescribing Information. Bedminster, NJ: Amarin Pharma, Inc.; December 2019. Available at: www.vascepa.com. Accessed November 2, 2020.
2. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2018;Nov 10:[Epub ahead of print].
3. Miller M, Stone NJ, Ballantyne C, et al. Triglycerides and cardiovascular disease: a scientific statement from the American Heart Association. *Circulation*. 2011; 123: 2292-2333.
4. Jellinger PS, Handelsman Y, Rosenblit PD, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Guidelines for management of dyslipidemia and prevention of cardiovascular disease. *Endocr Pract* 2017;23(Suppl 2):1-87.
5. Berglund L, Brunzell JD, Goldberg AC, et al. Evaluation and treatment of hypertriglyceridemia: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 2012;97(9):2969-2989.
6. Bhatt DL, Steg G, Miller M, et al. Cardiovascular risk reduction with icosapent ethyl for hypertriglyceridemia. *N Engl J Med*. 2019 Jan 1;380(1):11-22.
7. Skulas-Ray AC, Wilson PWF, Harris WS, et al. Omega-3 fatty acids for the management of hypertriglyceridemia: a science advisory from the American Heart Association. *Circulation*. 2019 Sep 17;140(12):e673-e691.
8. Manpuya WM, Cho L, Frid D, et al. Treatment strategies in patients with statin intolerance: the Cleveland Clinic experience. *American Heart Journal* 2013; 166(3):597-603.
9. Zhang H, Plutzky J, Skentzos S, et al. Discontinuation of statins in routine care settings. *Ann of Intern Med* 2013; 158(7):526-534.
10. Backes JM, Ruisinger JF, Gibson CA, et al. Statin-associated muscle symptoms—managing the highly intolerant. *J Clin Lipidol*. 2017;11:24-33. Available at: <https://www.acc.org/latest-in-cardiology/ten-points-to-remember/2017/05/03/10/43/statin-associated-muscle-symptoms>. Accessed December 5, 2019.
11. Thompson PD, Panza G, Zaleski A, et al. Statin-associated side effects. *JACC* 2016;67(20):2395-2410.
12. American Diabetes Association. 10. Cardiovascular disease and risk management: Standards of medical care in diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S111-S134.
13. Goff DC Jr, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014 Jun 24;129(25 Suppl 2):S49-73.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	01.22.20	02.20
1Q 2021 annual review: no significant changes; added redirection to generic icosapent ethyl for brand Vascepa requests; references reviewed and updated.	11.02.20	02.21
Removed redirection to generic icosapent ethyl per current rebate strategy.	10.07.21	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information.

This Clinical Policy is not intended to dictate to providers how to practice medicine, nor does it constitute a contract or guarantee regarding payment or results. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members.

This policy is the property of Envolve Pharmacy Solutions. Unauthorized copying, use, and distribution of this Policy or any information contained herein is strictly prohibited. By accessing this policy, you agree to be bound by the foregoing terms and conditions, in addition to the Site Use Agreement for Health Plans associated with Envolve Pharmacy Solutions.

©2020 Envolve Pharmacy Solutions. All rights reserved. All materials are exclusively owned by Envolve Pharmacy Solutions and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Envolve Pharmacy Solutions. You may not alter or remove any trademark, copyright or other notice contained herein.