

Clinical Policy: Anifrolumab-fnia (Saphnelo)

Reference Number: ERX.SPA.455

Effective Date: 12.01.21

Last Review Date: 11.21

Line of Business: Commercial, Medicaid

[Revision Log](#)

See **Important Reminder** at the end of this policy for important regulatory and legal information.

Description

Anifrolumab-fnia (Saphnelo[™]) is a type I interferon (IFN) receptor antagonist.

FDA Approved Indication(s)

Saphnelo is indicated for the treatment of adult patients with moderate to severe systemic lupus erythematosus (SLE) who are receiving standard therapy.

Limitation(s) of use: The efficacy of Saphnelo has not been evaluated in patients with severe active lupus nephritis or severe active central nervous system lupus. Use of Saphnelo is not recommended in these situations.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

Health plan approved formularies should be reviewed for all coverage determinations. Requirements to use preferred alternative agents apply only when such requirements align with the health plan approved formulary.

It is the policy of health plans affiliated with Envolve Pharmacy Solutions[™] that Saphnelo is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Systemic Lupus Erythematosus (must meet all):

1. Diagnosis of SLE;
2. Prescribed by or in consultation with a rheumatologist;
3. Age \geq 18 years;
4. Documentation confirms that member is positive for an SLE autoantibody (e.g., anti-nuclear antibody (ANA), anti-double-stranded DNA (anti-dsDNA), anti-Smith (anti-Sm), anti-ribonucleoprotein (anti-RNP), anti-Ro/SSA, anti-La/SSB, antiphospholipid antibody);
5. Prescribed in combination with standard therapy for SLE that includes one or more of the following agents, unless all agents are contraindicated: glucocorticoids (e.g., prednisone), antimalarial (e.g., hydroxychloroquine or chloroquine), non-biologic immunosuppressants (e.g., azathioprine, methotrexate, mycophenolate);
6. Dose does not exceed 300 mg every 4 weeks.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to ERX.PA.01 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions or member has previously met initial approval criteria;
2. Member is responding positively to therapy;

3. Prescribed in combination with standard therapy for SLE that includes one or more of the following agents, unless all agents are contraindicated: glucocorticoids (e.g., prednisone), antimalarial (e.g., hydroxychloroquine or chloroquine), non-biologic immunosuppressants (e.g., azathioprine, methotrexate, mycophenolate);
4. If request is for a dose increase, new dose does not exceed 300 mg every 4 weeks.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to ERX.PA.01 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – ERX.PA.01 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ANA: anti-nuclear antibody

Anti-dsDNA: anti-double-stranded DNA

Anti-Sm: anti-Smith

DNA: deoxyribonucleic acid

FDA: Food and Drug Administration

SLE: systemic lupus erythematosus

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria.

The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
glucocorticoids (e.g., prednisone)	Varies	Varies
antimalarial agents (e.g., hydroxychloroquine, chloroquine)	Varies	Varies
non-biologic immunosuppressants (e.g., azathioprine, methotrexate, mycophenolate)*	Varies	Varies

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

** For LN, cyclophosphamide is also an acceptable immunosuppressant.*

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): previous anaphylaxis with anifrolumab-fnia
- Boxed warning(s): none reported

Appendix D: Autoantibody Positive Versus Negative SLE

The pivotal clinical trials for Saphnelo enrolled patients with at least one of the following:

- Positive antinuclear antibody test at screening by immunofluorescent assay (IFA) at the central laboratory with titer \geq 1:80;
- Anti-dsDNA antibodies at screening elevated to above normal (including indeterminate), as per the central laboratory;
- Anti-Smith antibody at screening elevated to above normal as per the central laboratory

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
SLE	300 mg IV every 4 weeks	300 mg/4 weeks

VI. Product Availability

Single-dose vial: 300 mg/2 mL

VII. References

1. Saphnelo Prescribing Information. Wilmington, DE: AstraZeneca Pharmaceuticals LP; July 2021. Available at www.saphnelo.com. Accessed August 26, 2021.
2. Fanouriakis A, Kostopoulou M, Alunno A, et al. 2019 update of the EULAR recommendations for the management of systemic lupus erythematosus. *Ann Rheum Dis*. 2019;0:1–10. doi:10.1136/annrheumdis-2019-215089.
3. Petri M, Orbai AM, Alarcón GS, et al. Derivation and validation of the Systemic Lupus International Collaborating Clinics classification criteria for systemic lupus erythematosus. *Arthritis Rheum*. 2012; 64:2677.
4. Gordon C, Amissah-Arthur MB, Gayed M, et al. The British Society for Rheumatology guideline for the management of systemic lupus erythematosus in adults. *Rheumatology*. 2018;57:e1-e45. doi:10.1093/rheumatology/kex286.
5. Morand EF, Furie R, Tanaka Y, et al. Trial of Anifrolumab in Active Systemic Lupus Erythematosus. *N Engl J Med* 2020;382:211-21.
6. Furie R, Khamashta M, Merrill JT, et al. Anifrolumab, an Anti-Interferon-α Receptor Monoclonal Antibody, in Moderate-to-Severe Systemic Lupus Erythematosus. *Arthritis & Rheumatology* 2017; 69(2): 376-386.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	08.26.21	11.21

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information.

This Clinical Policy is not intended to dictate to providers how to practice medicine, nor does it constitute a contract or guarantee regarding payment or results. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members.

This policy is the property of Envolve Pharmacy Solutions. Unauthorized copying, use, and distribution of this Policy or any information contained herein is strictly prohibited. By accessing this policy, you agree to be bound by the foregoing terms and conditions, in addition to the Site Use Agreement for Health Plans associated with Envolve Pharmacy Solutions.

©2021 Envolve Pharmacy Solutions. All rights reserved. All materials are exclusively owned by Envolve Pharmacy Solutions and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Envolve Pharmacy Solutions. You may not alter or remove any trademark, copyright or other notice contained herein.